



TITLE:

Generation of a replication-competent simian-human immunodeficiency virus, the neutralization sensitivity of which can be enhanced in the presence of a small-molecule CD4 mimic(Dissertation_全文)

AUTHOR(S):

Otsuki, Hiroyuki

CITATION:

Otsuki, Hiroyuki. Generation of a replication-competent simian-human immunodeficiency virus, the neutralization sensitivity of which can be enhanced in the presence of a small-molecule CD4 mimic. 京都大学, 2014, 博士(医科学)

ISSUE DATE:

2014-03-24

URL:

<https://doi.org/10.14989/doctor.k18186>

RIGHT:

許諾条件により本文は2014-09-12に公開; This is an author manuscript that has been accepted for publication in Journal of General Virology, copyright Society for General Microbiology, but has not been copy-edited, formatted or proofed. Cite this article as appearing in Journal of General Virology. This version of the manuscript may not be duplicated or reproduced, other than for personal use or within the rule of 'Fair Use of Copyrighted Materials' (section 17, Title 17, US Code), without permission from the copyright owner, Society for General Microbiology. The Society for General Microbiology disclaims any responsibility or liability for errors or omissions in this version of the manuscript or in any version derived from it by any other parties. The final copy-edited, published article, which is the version of record, can be found at <http://vir.sgmjournals.org>, and is freely available without a ...

主論文Web公開版

This is an author manuscript that has been accepted for publication in Journal of General Virology, copyright Society for General Microbiology, but has not been copy-edited, formatted or proofed. Cite this article as appearing in Journal of General Virology. This version of the manuscript may not be duplicated or reproduced, other than for personal use or within the rule of 'Fair Use of Copyrighted Materials' (section 17, Title 17, US Code), without permission from the copyright owner, Society for General Microbiology. The Society for General Microbiology disclaims any responsibility or liability for errors or omissions in this version of the manuscript or in any version derived from it by any other parties. The final copy-edited, published article, which is the version of record, can be found at <http://vir.sgmjournals.org>, and is freely available without a subscription.

Title page

- Generation of a replication-competent simian–human immunodeficiency virus, the neutralisation sensitivity of which can be enhanced in the presence of a small molecule CD4 mimic
- The Contents Category for the paper, Short communications
- Short running title: SHIV with conditional neutralisation sensitivity
- The names of the authors, Hiroyuki Otsuki¹, Tomoyuki Miura¹, Chie Hashimoto², Tetsuo Narumi², Hirokazu Tamamura², Kazuhisa Yoshimura³, Shuzo Matsushita⁴, and Tatsuhiko Igarashi^{1#}
- The name and address of the laboratories where the work was done,
¹Laboratory of Primate Model, Experimental Research Center for Infectious Diseases, Institute for Virus Research, Kyoto University, Kyoto 606-8507, Japan,
²Department of Medicinal Chemistry, Institute of Biomaterials and Bioengineering, Tokyo Medical and Dental University, Tokyo 101-0062, Japan, ³AIDS Research Center, National Institute of Infectious Diseases, Tokyo 162-8640, Japan,
⁴Division of Clinical Retrovirology and Infectious Diseases, Center for AIDS Research, Kumamoto University, Kumamoto 860-0811, Japan
- An email address and telephone and fax numbers for the corresponding author,
tigarash@virus.kyoto-u.ac.jp, E-mail
81 75 751 3982, Phone
81 75 761 9335, Fax
- The word count of the summary and the main text,
Summary, 143 words
Main text, 2451 words
Figure legends, 225 words

Summary

Simian-human immunodeficiency virus (SHIV) carrying the envelope from the clade B clinical HIV-1 isolate HIV-1 MNA, designated SHIV MNA, was generated through intracellular homologous recombination. SHIV MNA inherited biological properties from the parental HIV-1, including CCR5 co-receptor preference, resistance to neutralisation by the anti-V3 loop monoclonal antibody KD-247, and loss of resistance in the presence of the CD4-mimic small molecule YYA-021. SHIV MNA showed productive replication in rhesus macaque peripheral blood mononuclear cells.

Experimental infection of a rhesus macaque with SHIV MNA caused a transient but high titre of plasma viral RNA and a moderate antibody response. Immunoglobulin in the plasma at 24 weeks post-infection was capable of neutralising SHIV MNA in the presence but not in the absence of YYA-021.

SHIV MNA could serve a model for development of novel therapeutic interventions based on CD4-mimic-mediated conversion of Env susceptible to antibody neutralisation.

Text

Control of primate lentiviral infection by antibodies directed against viral envelope protein is theoretically feasible. This was confirmed by the successful protection of macaque monkeys from challenge inoculation with simian-human immunodeficiency virus (SHIV) carrying an envelope protein (Env). Env was derived from a laboratory strain of human immunodeficiency virus type 1 (HIV-1) through the passive immunisation of neutralising monoclonal antibodies directed against HIV-1 (Mascola *et al.*, 2000; Nishimura *et al.*, 2003). This neutralisation is consistent with the results normally seen in cell culture systems.

Clinical isolates of HIV-1, which have not been subjected to extensive passage in T-cell lines, on the other hand, are generally resistant to antibody-mediated neutralisation (Moore *et al.*, 1995). It has been shown that virus in infected individuals is under selective pressure to develop a variety of means to evade attack by neutralising antibodies, including sequence variation, glycosylation, tertiary structural shielding formed by the Env trimer, and the rapid kinetics of conformational changes of Env, which affect fusion between the viral envelope and the plasma membrane of target cells

(Kong & Sattentau, 2012). Although four major neutralising epitopes have been identified in the HIV-1 Env; *i.e.* the V1/V2 loop, the glycan-V3 site and CD4-binding site of gp120, and the membrane-proximal external region (MPER) of gp41, few reports of antibodies directed against these epitopes capable of neutralising a broad range of isolates have been published, for reasons that are as yet unclear (Kwong & Mascola, 2012). High titres of antibodies directed against the V3 loop are elicited in individuals during the early phase of HIV-1 infection, but these are incapable of neutralising the virus because the epitope in functional Env trimer is likely shielded from the antibody (Davis *et al.*, 2009b). Therefore, it is necessary to develop a means of rendering these epitopes accessible to the antibodies, to make antibody-mediated suppression of HIV-1 a valid therapeutic option.

It has been reported that neutralisation mediated by antibodies directed against the V3 loop (Lusso *et al.*, 2005) or CD4-induced epitope (CD4i) (Thali *et al.*, 1993) can be enhanced in the presence of soluble CD4 (sCD4). It is known that the interaction of Env with sCD4 drives a conformational change of the viral protein and makes the cryptic/occult epitopes accessible to these antibodies (Wyatt *et al.*, 1998). Small

molecules that emulate sCD4 for its interaction and subsequent induction of conformational change of Env may be employed to intensify antibody-mediated interventions against HIV-1 infection. Compounds with the above-mentioned properties; *i.e.* NBD-556 and NBD-557, have been reported previously (Zhao *et al.*, 2005). NBD-556 has been shown in cell culture to interact with the CD4-binding pocket to induce a conformational change in gp120 (Madani *et al.*, 2008) and enhance exposure of the Env of primary HIV-1 isolates to neutralising epitopes (Yoshimura *et al.*, 2010).

The present study was performed to evaluate small molecule CD4-mimic-based enhancement of antibody-mediated virus neutralisation, in the context of virus infection *in vivo*. The simian-human immunodeficiency virus (SHIV)/macaque monkey model of AIDS is particularly suitable for such studies, as SHIV carries the HIV-1 Env and the neutralisation sensitivity of SHIV is comparable to that of the parental HIV-1 (Shibata & Adachi, 1992).

As NBD-556, unlike sCD4, inhibits infection with select HIV-1 isolates (Yoshimura *et al.*, 2010), we generated a new SHIV strain carrying Env, the sensitivity of which to antibody-mediated neutralisation is enhanced in the presence of a CD4 mimic. An

HIV-1 isolate, MNA, previously designated primary isolate HIV-1 Pt.3, was used as the source of Env, as the viral protein has been reported to interact with NBD-556 (Yoshimura *et al.*, 2010). While the virus belongs to a distinct subset of HIV-1 isolates, as mentioned above, it has also been reported to utilise the CCR5 molecule to gain entry into target cells, a property that is shared by the majority of HIV-1 strains (Yoshimura *et al.*, 2010). A monoclonal antibody directed against the tip of the V3 loop (GPGR), KD-247 (Eda *et al.*, 2006), was employed to assess this concept, as HIV-1 MNA was resistant to KD-247-mediated neutralisation, despite carrying the exact epitope sequence in the tip of V3 loop, and was converted to being sensitive to the antibody by NBD-556 in a dose-dependent manner (Yoshimura *et al.*, 2010).

First, we reproduced the results of Yoshimura *et al.* using a neutralisation assay employing TZM-bl cells (Platt *et al.*, 1998), obtained from the NIH AIDS Reagent program (Fig. S1). The virus was resistant to KD-247, as described previously, and required almost 50 $\mu\text{g/mL}$ of the antibody to achieve 50% neutralisation in our assay. The observed resistance was abrogated in the presence of 2 μM of NBD-556. However, 50% neutralisation was achieved in the presence of ~ 0.1 $\mu\text{g/mL}$ of KD-247,

corresponding to 1/500 of the amount of the antibody to achieve the same degree of neutralisation in the absence of the CD4 mimic.

With reproduction of the properties of HIV-1 MNA Env, we generated an SHIV strain carrying Env through intracellular homologous recombination, as described previously (Fujita *et al.*, 2013) with minor modifications (Fig. S2). DNA fragments representing the 5' and 3' ends of the SHIV genome (fragments I and II, respectively) were amplified by PCR from the proviral DNA plasmid SHIV KS661. A DNA fragment containing *env* (fragment III) was amplified from complementary DNA (cDNA) of the HIV-1 MNA genome, which was prepared from virus particles (virion-associated RNA) in the culture supernatant of PM1/CCR5 cells (Yusa *et al.*, 2005) infected with the virus. The PCR primers used are listed in Table S1. Using a FuGENE HD transfection reagent, lipofection was performed on the C8166-CCR5 cells (Shimizu *et al.*, 2006) to co-transfect them with 0.2 µg of DNA. A cytopathic effect presumably caused by the emerged recombinant virus was observed on day 13 post-transfection. The emerged virus, designated SHIV MNA, carried the entire gp120 and three quarters of gp41 from HIV-1 MNA Env (Fig. 1a). The rest of Env was from

127 SHIV KS661, the Env of which was derived from HIV-1 89.6 (Shinohara *et al.*, 1999).
128 The CD4 binding site, and the regions and elements that reportedly interact with
129 NBD-556 (Madani *et al.*, 2008; Yoshimura *et al.*, 2010), are preserved in SHIV MNA
130 Env (Fig. S3). The virus was replication-competent in PM1/CCR5 cells (data not
131 shown).

132 As HIV-1 MNA was suggested to be a CCR5-utilising virus, we were intrigued
133 whether SHIV MNA inherited the trait from the parental virus. We subjected SHIV
134 MNA and the parental HIV-1 MNA to co-receptor usage assay as described previously
135 (Nishimura *et al.*, 2010), with minor modifications (Fig. S4). As expected, SHIV MNA
136 was shown to utilise CCR5 as an entry co-receptor.

137 We next assessed the neutralisation profiles of SHIV MNA in comparison with the
138 parental HIV-1 MNA, as described previously (Li *et al.*, 2005; Wei *et al.*, 2002). Both
139 SHIV MNA and HIV-1 MNA showed essentially no neutralisation by KD-247 up to 25
140 µg/mL and 50% neutralisation was achieved at 50 µg/mL (Fig. 1b). As the CD4 mimic,
141 we employed YYA-021, a compound generated and characterised through studies
142 concerning the structure-activity relationships of small molecules (Narumi *et al.*, 2013;

143 Narumi *et al.*, 2011; Narumi *et al.*, 2010; Yamada *et al.*, 2010). The compound was
144 shown to be slightly less potent but to exhibit substantially lower toxicity than
145 NBD-556, and was therefore a suitable choice for our purposes in future studies in
146 animal models. SHIV MNA was resistant to neutralisation by YYA-021 at all
147 concentrations examined, except 25 and 50 μ M, and showed a neutralisation profile
148 almost identical to that of HIV-1 MNA (Fig. 1c). To further characterise the biological
149 properties of SHIV MNA Env, a set of entry assays was conducted (Fig. S5). The *env*
150 genes cloned from SHIV MNA and HIV-1 MNA, were utilised to generate
151 pseudo-typed viruses. These pseudotypes were inoculated into TZM-bl cells in the
152 presence of increasing amounts of NBD-556, YYA-021 or soluble CD4. A control
153 group was derived from another virus preparation pseudotyped with A-MLV Env
154 (Landau *et al.*, 1991). When the efficiency of entry was defined by intracellular
155 luciferase activities, virtually no difference was observed between Envs of SHIV MNA
156 and the parental HIV-1. Thus SHIV MNA Env replicated in C8166-CCR5 cells retained
157 sensitivity to small molecule CD4 mimics and soluble CD4 comparable to that of
158 HIV-1 MNA.

We next examined whether the synergistic neutralisation of HIV-1 MNA by KD-247 antibody in the presence of NBD-556 (Yoshimura *et al.*, 2010) would be reproduced when CD4 mimic was substituted by YYA-021. The synergistic neutralisation effect of KD-247 and YYA-021 was reproduced in our experiments (Fig. 1d). At 50 µg/mL, KD-247 barely achieved 50% neutralisation of HIV-1 MNA but resulted in 50% neutralisation at < 0.05 µg/mL in the presence of 20 µM of YYA-021.

Finally, to examine whether these two agents neutralise SHIV MNA in the same manner as the parental HIV-1, we conducted a neutralisation assay with KD-247 in the presence of increasing amounts of YYA-021 (0, 5, 10, 20 and 40 µM) (Fig. 1e). The neutralisation curve of KD-247 against SHIV MNA showed an upward shift as the concentration of YYA-021 increased (Fig. 1e), similar to the observations with HIV-1 (Fig. 1d), indicating augmentation of neutralisation, and complete neutralisation of both viruses was achieved at 20 µM YYA-021 (Fig. 1d and e). Based on these results, we concluded that the neutralisation profile of SHIV MNA was comparable to that of HIV-1 MNA.

Reproduction of the neutralisation characteristics of HIV-1 MNA in the newly

generated SHIV prompted us to assess the ability of SHIV MNA to replicate in monkey cells. SHIV MNA, along with SIV239 and SHIV KS661, were normalised with infectious titres and inoculated into rhesus macaque peripheral blood mononuclear cell (PBMC) preparations from four animals, as described previously (Fujita *et al.*, 2013) (Fig. 2a). SHIV KS661, a CXCR4-utilising virus, replicated to the highest titres of all the viruses in all PBMC preparations. Compared to SHIV KS661, SIV239 replicated to lower titres. Under these experimental conditions, SHIV MNA showed productive replication in the cells with similar replication kinetics and peak titres to SIV239. Based on these results, we concluded that SHIV MNA was replication-competent in primary monkey lymphocytes.

Productive replication of SHIV MNA in monkey PBMC justified experimental infection of the virus *in vivo*. We inoculated 1.75×10^5 TCID₅₀ of SHIV MNA intravenously into a rhesus macaque and monitored plasma viral RNA burden and circulating CD4⁺ T-lymphocyte levels. Plasma viral RNA burden reached a peak of 5.6×10^6 copies/mL at 1 week post-infection (wpi), and declined rapidly thereafter reaching low levels of detection at 7 wpi (around 2.8×10^2 copies/mL). Circulating CD4⁺

191 T-cell numbers showed a transient decrease around 1 wpi, rebounded around 3 wpi and
192 stabilised around 70% of the pre-infection level from 4 wpi. During the period of
193 observation, the animal developed no obvious clinical manifestations related to
194 lentivirus infection.

195 As SHIV MNA replicated *in vivo* without depleting helper T-cells, it was expected
196 that the animal mounted an anti-viral immune reaction. The production of antibody
197 directed against Env was assessed by western immunoblotting, as described previously
198 (Igarashi *et al.*, 1999). Purified Env protein (Advanced Biotechnologies Inc. Md.
199 U.S.A.) was used as the antigen (Fig. 3a). Anti-Env antibody was detected at 3 wpi and
200 the level of antibody—judged by the intensity of the band—increased gradually with
201 time.

202 We next examined whether the animal generated neutralising antibodies against
203 SHIV MNA. Because plasma samples from this specimen exhibited high background
204 activity, immunoglobulin G (IgG) was purified from these samples collected on day 0
205 and in week 24 post-infection using protein G spin columns (GE healthcare Japan.
206 Tokyo. Japan). While the IgG from day 0 exhibited no neutralising activity (Fig. 3b), as

expected, the immunoglobulin collected at 24 wpi neutralised SHIV MNA, although a concentration $> 100 \mu\text{g/mL}$ was required to suppress replication of 100 TCID₅₀ of the input virus (Fig. 3c).

We examined whether the observed marginal neutralisation by the antibody could be enhanced by the presence of YYA-021. Upon addition of YYA-021 in the assay system, SHIV MNA became sensitive to IgG obtained at 24 wpi (Fig. 3c), while no enhancement was identified from day 0 (Fig. 3b).

In this study, we generated a replication-competent SHIV MNA strain carrying an Env resistant to the monoclonal neutralising antibody KD-247 but conditionally sensitive in the presence of the CD4 mimic YYA-021. As the observed neutralisation characteristics were identical to those of HIV-1 MNA, which contributed the majority of the Env sequence to the chimera, the utility of the CD4 mimic as a means of enhancing antibody-mediated virus neutralisation should be assessed in the context of infection *in vivo*. This concept could be tested during the acute phase of SHIV MNA infection, during which the virus undergoes substantial replication. To examine the feasibility of CD4-mimic-mediated enhancement of virus neutralisation in the context

223 of chronic infection, the conditions under which this type of intervention should be
224 applied to HIV-1-infected patients in a clinical setting, the virus must be modified to
225 sustain productive replication for a longer period. SHIV MNA in the present form does
226 not fulfil this requirement. It is possible that animal-to-animal passage could increase
227 the fitness of the virus in monkeys.

228 This study demonstrated that a CD4 mimic could modulate viral Env protein to be
229 more susceptible to neutralisation by less potent antibodies generated in the context of
230 infection. During the early phase of infection, patients mount high titres of
231 non-neutralising antibodies directed against the V3 loop (Davis *et al.*, 2009a). Patients
232 with HIV-1 clade C generate anti-Env antibodies, including anti-CD4i antibodies, with
233 poor neutralising activity against recent infection (Gray *et al.*, 2007). It is possible that
234 the CD4 mimic YYA-021 causes a conformational change in SHIV MNA Env, which
235 renders sequestered epitope(s) accessible to potentially neutralising IgG, such as the V3
236 loop and CD4i.

237 The current study extended the previous study by Yoshimura *et al.* and used HIV-1
238 MNA belonging to clade B to generate a new SHIV strain carrying Env. The

neutralisation sensitivity of this strain is characteristically augmented in the presence of a small molecule CD4 mimic. Similar observations by Decker *et al.* show that infections of a wide range of HIV-1 strains of multiple clades or circulating recombinant forms elicits high titres of anti-CD4i antibodies, These anti-CD4i antibodies neutralise viruses as divergent as HIV-2 in the presence of soluble CD4 (Decker *et al.*, 2005). Taking these observations into account, small molecule CD4 mimics such as YYA-021 could potentially enhance the neutralisation activity of the antibodies directed against autologous viruses belonging not only to clade B but also to multiple HIV-1 strains of various clades, and possibly even HIV-2. Our results pave the way for a novel therapeutic intervention based on administration of CD4 mimics to patients with HIV to facilitate control of the virus by their own antibodies.

250

Acknowledgements

251 Thanks should be given to: Drs. Julie Strizki and Paul Zavodny of the Schering-Plough
252 Research Institute, Kenilworth, NJ. U.S.A. for providing AD101; the NIH AIDS
253 Research & Reference Reagent Program for providing TZM-bl cells, SV-A-MLV-env,
254 4G10 and soluble CD4; the Chemo-Sero-Therapeutic Research Institute (Kaketsuken)
255 for providing MAb KD-247; former and current members of the Igarashi laboratory for
256 discussion and support. This work was supported by a Research on HIV/AIDS grant
257 (H22-AIDS Research-007 and H24-AIDS Research-008) from The Ministry of Health,
258 Labor and Welfare of Japan and by a Grant-in-Aid for Scientific Research (B)
259 (23300156) from the Japan Society for the Promotion of Science.

260

- 261 **Davis, K. L., Bibollet-Ruche, F., Li, H., Decker, J. M., Kutsch, O., Morris, L.,**
 262 **Salomon, A., Pinter, A., Hoxie, J. A., Hahn, B. H., Kwong, P. D. & Shaw, G. M.**
 263 **(2009a).** Human immunodeficiency virus type 2 (HIV-2)/HIV-1 envelope chimeras
 264 detect high titers of broadly reactive HIV-1 V3-specific antibodies in human plasma.
 265 *Journal of virology* **83**, 1240-1259.
- 266 **Davis, K. L., Gray, E. S., Moore, P. L., Decker, J. M., Salomon, A., Montefiori, D.**
 267 **C., Graham, B. S., Keefer, M. C., Pinter, A., Morris, L., Hahn, B. H. & Shaw, G.**
 268 **M. (2009b).** High titer HIV-1 V3-specific antibodies with broad reactivity but low
 269 neutralizing potency in acute infection and following vaccination. *Virology* **387**,
 270 414-426.
- 271 **Decker, J. M., Bibollet-Ruche, F., Wei, X., Wang, S., Levy, D. N., Wang, W.,**
 272 **Delaporte, E., Peeters, M., Derdeyn, C. A., Allen, S., Hunter, E., Saag, M. S.,**
 273 **Hoxie, J. A., Hahn, B. H., Kwong, P. D., Robinson, J. E. & Shaw, G. M. (2005).**
 274 Antigenic conservation and immunogenicity of the HIV coreceptor binding site. *The*
 275 *Journal of experimental medicine* **201**, 1407-1419.
- 276 **Donzella, G. A., Schols, D., Lin, S. W., Este, J. A., Nagashima, K. A., Maddon, P. J.,**
 277 **Allaway, G. P., Sakmar, T. P., Henson, G., De Clercq, E. & Moore, J. P. (1998).**
 278 AMD3100, a small molecule inhibitor of HIV-1 entry via the CXCR4 co-receptor.
 279 *Nature medicine* **4**, 72-77.
- 280 **Eda, Y., Takizawa, M., Murakami, T., Maeda, H., Kimachi, K., Yonemura, H.,**
 281 **Koyanagi, S., Shiosaki, K., Higuchi, H., Makizumi, K., Nakashima, T., Osatomi,**
 282 **K., Tokiyoshi, S., Matsushita, S., Yamamoto, N. & Honda, M. (2006).** Sequential
 283 immunization with V3 peptides from primary human immunodeficiency virus type 1
 284 produces cross-neutralizing antibodies against primary isolates with a matching
 285 narrow-neutralization sequence motif. *Journal of virology* **80**, 5552-5562.
- 286 **Fujita, Y., Otsuki, H., Watanabe, Y., Yasui, M., Kobayashi, T., Miura, T. &**
 287 **Igarashi, T. (2013).** Generation of a replication-competent chimeric simian-human
 288 immunodeficiency virus carrying env from subtype C clinical isolate through
 289 intracellular homologous recombination. *Virology* **436**, 100-111.
- 290 **Gray, E. S., Moore, P. L., Choge, I. A., Decker, J. M., Bibollet-Ruche, F., Li, H.,**
 291 **Leseka, N., Treurnicht, F., Mlisana, K., Shaw, G. M., Karim, S. S., Williamson,**

- C., Morris, L. & Team, C. S. (2007). Neutralizing antibody responses in acute human immunodeficiency virus type 1 subtype C infection. *Journal of virology* **81**, 6187-6196.
- Igarashi, T., Endo, Y., Englund, G., Sadjadpour, R., Matano, T., Buckler, C., Buckler-White, A., Plishka, R., Theodore, T., Shibata, R. & Martin, M. (1999). Emergence of a highly pathogenic simian/human immunodeficiency virus in a rhesus macaque treated with anti-CD8 mAb during a primary infection with a nonpathogenic virus. *Proceedings of the National Academy of Sciences of the United States of America* **96**, 14049-14054.
- Kong, L. & Sattentau, Q. J. (2012). Antigenicity and Immunogenicity in HIV-1 Antibody-Based Vaccine Design. *J AIDS Clin Res* **S8**, 3.
- Kwong, P. D. & Mascola, J. R. (2012). Human antibodies that neutralize HIV-1: identification, structures, and B cell ontogenies. *Immunity* **37**, 412-425.
- Landau, N. R., Page, K. A. & Littman, D. R. (1991). Pseudotyping with human T-cell leukemia virus type I broadens the human immunodeficiency virus host range. *Journal of virology* **65**, 162-169.
- Li, M., Gao, F., Mascola, J. R., Stamatatos, L., Polonis, V. R., Koutsoukos, M., Voss, G., Goepfert, P., Gilbert, P., Greene, K. M., Bilska, M., Kothe, D. L., Salazar-Gonzalez, J. F., Wei, X., Decker, J. M., Hahn, B. H. & Montefiori, D. C. (2005). Human immunodeficiency virus type 1 env clones from acute and early subtype B infections for standardized assessments of vaccine-elicited neutralizing antibodies. *Journal of virology* **79**, 10108-10125.
- Lusso, P., Earl, P. L., Sironi, F., Santoro, F., Ripamonti, C., Scarlatti, G., Longhi, R., Berger, E. A. & Burastero, S. E. (2005). Cryptic nature of a conserved, CD4-inducible V3 loop neutralization epitope in the native envelope glycoprotein oligomer of CCR5-restricted, but not CXCR4-using, primary human immunodeficiency virus type 1 strains. *Journal of virology* **79**, 6957-6968.
- Madani, N., Schon, A., Princiotta, A. M., Lalonde, J. M., Courter, J. R., Soeta, T., Ng, D., Wang, L., Brower, E. T., Xiang, S. H., Kwon, Y. D., Huang, C. C., Wyatt, R., Kwong, P. D., Freire, E., Smith, A. B., 3rd & Sodroski, J. (2008). Small-molecule CD4 mimics interact with a highly conserved pocket on HIV-1 gp120. *Structure* **16**, 1689-1701.
- Mascola, J. R., Stiegler, G., VanCott, T. C., Katinger, H., Carpenter, C. B., Hanson,

325 **C. E., Beary, H., Hayes, D., Frankel, S. S., Birx, D. L. & Lewis, M. G. (2000).**
 326 Protection of macaques against vaginal transmission of a pathogenic HIV-1/SIV
 327 chimeric virus by passive infusion of neutralizing antibodies. *Nature medicine* **6**,
 328 207-210.

329 **Moore, J. P., Cao, Y., Qing, L., Sattentau, Q. J., Pyati, J., Koduri, R., Robinson, J.,**
 330 **Barbas, C. F., 3rd, Burton, D. R. & Ho, D. D. (1995).** Primary isolates of human
 331 immunodeficiency virus type 1 are relatively resistant to neutralization by
 332 monoclonal antibodies to gp120, and their neutralization is not predicted by studies
 333 with monomeric gp120. *Journal of virology* **69**, 101-109.

334 **Narumi, T., Arai, H., Yoshimura, K., Harada, S., Hirota, Y., Ohashi, N.,**
 335 **Hashimoto, C., Nomura, W., Matsushita, S. & Tamamura, H. (2013).** CD4
 336 mimics as HIV entry inhibitors: lead optimization studies of the aromatic
 337 substituents. *Bioorganic & medicinal chemistry* **21**, 2518-2526.

338 **Narumi, T., Arai, H., Yoshimura, K., Harada, S., Nomura, W., Matsushita, S. &**
 339 **Tamamura, H. (2011).** Small molecular CD4 mimics as HIV entry inhibitors.
 340 *Bioorganic & medicinal chemistry* **19**, 6735-6742.

341 **Narumi, T., Ochiai, C., Yoshimura, K., Harada, S., Tanaka, T., Nomura, W., Arai,**
 342 **H., Ozaki, T., Ohashi, N., Matsushita, S. & Tamamura, H. (2010).** CD4 mimics
 343 targeting the HIV entry mechanism and their hybrid molecules with a CXCR4
 344 antagonist. *Bioorganic & medicinal chemistry letters* **20**, 5853-5858.

345 **Nishimura, Y., Igarashi, T., Haigwood, N. L., Sadjadpour, R., Donau, O. K.,**
 346 **Buckler, C., Plishka, R. J., Buckler-White, A. & Martin, M. A. (2003).** Transfer
 347 of neutralizing IgG to macaques 6 h but not 24 h after SHIV infection confers
 348 sterilizing protection: implications for HIV-1 vaccine development. *Proceedings of*
 349 *the National Academy of Sciences of the United States of America* **100**, 15131-15136.

350 **Nishimura, Y., Shingai, M., Willey, R., Sadjadpour, R., Lee, W. R., Brown, C. R.,**
 351 **Brenchley, J. M., Buckler-White, A., Petros, R., Eckhaus, M., Hoffman, V.,**
 352 **Igarashi, T. & Martin, M. A. (2010).** Generation of the pathogenic R5-tropic
 353 simian/human immunodeficiency virus SHIVAD8 by serial passaging in rhesus
 354 macaques. *Journal of virology* **84**, 4769-4781.

355 **Platt, E. J., Wehrly, K., Kuhmann, S. E., Chesebro, B. & Kabat, D. (1998).** Effects
 356 of CCR5 and CD4 cell surface concentrations on infections by macrophagetropic
 357 isolates of human immunodeficiency virus type 1. *Journal of virology* **72**,

2855-2864.

Shibata, R. & Adachi, A. (1992). SIV/HIV recombinants and their use in studying biological properties. *AIDS research and human retroviruses* **8**, 403-409.

Shimizu, Y., Okoba, M., Yamazaki, N., Goto, Y., Miura, T., Hayami, M., Hoshino, H. & Haga, T. (2006). Construction and in vitro characterization of a chimeric simian and human immunodeficiency virus with the RANTES gene. *Microbes and infection / Institut Pasteur* **8**, 105-113.

Shinohara, K., Sakai, K., Ando, S., Ami, Y., Yoshino, N., Takahashi, E., Someya, K., Suzaki, Y., Nakasone, T., Sasaki, Y., Kaizu, M., Lu, Y. & Honda, M. (1999). A highly pathogenic simian/human immunodeficiency virus with genetic changes in cynomolgus monkey. *The Journal of general virology* **80** (Pt 5), 1231-1240.

Thali, M., Moore, J. P., Furman, C., Charles, M., Ho, D. D., Robinson, J. & Sodroski, J. (1993). Characterization of conserved human immunodeficiency virus type 1 gp120 neutralization epitopes exposed upon gp120-CD4 binding. *Journal of virology* **67**, 3978-3988.

Trkola, A., Kuhmann, S. E., Strizki, J. M., Maxwell, E., Ketas, T., Morgan, T., Pugach, P., Xu, S., Wojcik, L., Tagat, J., Palani, A., Shapiro, S., Clader, J. W., McCombie, S., Reyes, G. R., Baroudy, B. M. & Moore, J. P. (2002). HIV-1 escape from a small molecule, CCR5-specific entry inhibitor does not involve CXCR4 use. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 395-400.

von Brunn, A., Brand, M., Reichhuber, C., Morys-Wortmann, C., Deinhardt, F. & Schodel, F. (1993). Principal neutralizing domain of HIV-1 is highly immunogenic when expressed on the surface of hepatitis B core particles. *Vaccine* **11**, 817-824.

Wei, X., Decker, J. M., Liu, H., Zhang, Z., Arani, R. B., Kilby, J. M., Saag, M. S., Wu, X., Shaw, G. M. & Kappes, J. C. (2002). Emergence of Resistant Human Immunodeficiency Virus Type 1 in Patients Receiving Fusion Inhibitor (T-20) Monotherapy. *Antimicrobial Agents and Chemotherapy* **46**, 1896-1905.

Wyatt, R., Kwong, P. D., Desjardins, E., Sweet, R. W., Robinson, J., Hendrickson, W. A. & Sodroski, J. G. (1998). The antigenic structure of the HIV gp120 envelope glycoprotein. *Nature* **393**, 705-711.

Yamada, Y., Ochiai, C., Yoshimura, K., Tanaka, T., Ohashi, N., Narumi, T., Nomura, W., Harada, S., Matsushita, S. & Tamamura, H. (2010). CD4 mimics

targeting the mechanism of HIV entry. *Bioorganic & medicinal chemistry letters* **20**, 354-358.

Yoshimura, K., Harada, S., Shibata, J., Hatada, M., Yamada, Y., Ochiai, C., Tamamura, H. & Matsushita, S. (2010). Enhanced exposure of human immunodeficiency virus type 1 primary isolate neutralization epitopes through binding of CD4 mimetic compounds. *Journal of virology* **84**, 7558-7568.

Yusa, K., Maeda, Y., Fujioka, A., Monde, K. & Harada, S. (2005). Isolation of TAK-779-resistant HIV-1 from an R5 HIV-1 GP120 V3 loop library. *The Journal of biological chemistry* **280**, 30083-30090.

Zhao, Q., Ma, L., Jiang, S., Lu, H., Liu, S., He, Y., Strick, N., Neamati, N. & Debnath, A. K. (2005). Identification of N-phenyl-N'-(2,2,6,6-tetramethyl-piperidin-4-yl)-oxalamides as a new class of HIV-1 entry inhibitors that prevent gp120 binding to CD4. *Virology* **339**, 213-225.

Figure legends

Figure 1. Genomic organisation (a) and neutralisation sensitivity (b – e) of SHIV MNA.

(a) Grey boxes represent genes derived from SIV239, open boxes those from HIV-1 89.6 and filled dark grey boxes those from HIV-1 MNA. (b – e) Percentage neutralisation was calculated as follows: % neutralisation = $100 \times \{1 - (\text{RLU.N} - \text{RLU.B})/(\text{RLU.V} - \text{RLU.B})\}$. RLU, relative luciferase units; RLU.N, RLU in wells with cells, virus and KD-247 and/or YYA-021; RLU.V, RLU in wells with cells and virus; RLU.B, RLU in wells with cells.

Figure 2. Replication of SHIV MNA in rhesus macaque PBMCs (a) and *in vivo* (b). (a) Multiplicity of infection was adjusted to 0.01 (TCID₅₀/cell). (b) Experimental infection of a rhesus macaque with SHIV MNA. SHIV MNA (1.75×10^5 TCID₅₀) was intravenously inoculated into a rhesus macaque, and the plasma viral RNA burden (filled circles) and circulating CD4⁺ T-lymphocytes (open triangles) were monitored.

Figure 3. Antibody induced against SHIV MNA.

423 (a) The anti-HIV-1 gp120 antibody response was assessed by immunoblotting with
424 plasma samples collected at the indicated times. An anti-HIV-1 V3 monoclonal
425 antibody 4G10 ascites (1:100) (von Brunn *et al.*, 1993), obtained from the NIH AIDS
426 Reagent program, was used as a positive control (lane anti-V3). (b and c) Neutralisation
427 of SHIV MNA with IgG purified from plasma of the infected rhesus macaque (day 0
428 and 24 wpi) with/without YYA-021 (20 μ M).

1 Table S1. PCR primers.

Primer	Sequence	Position (nt [†])
Fragment I		
SIVU3Not-F	5'-atgcggccgctggaagggatttattacagtgaag-3'	1 – 25 [*]
Preenv-R	5'-aaagagcagaagacgagtggcaa-3'	6204 – 6226 [#]
Fragment II		
SHenv5.5F	5'-tcataatgatagtaggaggc-3'	8278 – 8297 [#]
SIVU5Eco-R	5'-tgcagaattctgctagggattttcctgcttcggtt-3'	10255 – 10279 [*]
Fragment III		
HIV-1vpr-F	5'-agatggaacaagccccagaaga-3'	5557 – 5578 [#]
SHenv6R	5'-gctgaagaggcacaggctccgc-3'	8525 – 8504 [#]

2 [†], Nucleotide positions of PCR primers were numbered relative to the SIV239 (*,

3 GenBank Accession No. M33262) or HXB2 (#, GenBank Accession No. K03455)

4 genome sequences.

Figure S1. Enhanced neutralisation of HIV-1 MNA by KD-247 in the presence of

NBD-556.

100 TCID₅₀ of HIV-1 MNA was pre-incubated with increasing amounts of KD-247

with/without 2 µM of NBD-556 at 37°C for 60 min, followed by inoculation into 5×10³

TZM-bl cells. The cells were lysed at 48 h post-infection and luciferase activity was

measured. The percentage of neutralisation was measured as RLU reduction relative to

virus control wells.

Figure S2. Genomic organisation of SHIV KS661 and HIV-1 MNA and PCR fragments

employed for preparation of DNA fragments for generation of SHIV MNA.

Colour-coded boxes represent genes derived from the following viruses: grey boxes,

SIV239; open boxes, HIV-1 89.6; grey boxes, HIV-1 MNA. SHIV KS661 carries *tat*,

rev, *vpu*, and *env* genes from subtype B HIV-1 89.6. Broad lines represent PCR

fragments that were amplified using the primers indicated by arrows (A – F).

20 Figure S3. Deduced amino acid sequence alignment of Env from HIV-1 MNA, SHIV
21 MNA, and SHIV KS661.
22 (*) = Amino acids that form part of the CD4 binding site.
23 (†) = Regions/elements that are reported to interact with NBD-556 (Madani *et al.*, 2008;
24 Yoshimura *et al.*, 2010).
25 Parts of SHIV MNA Env, that were putatively derived from HIV-1 MNA or SHIV
26 KS661 are respectively color-coded as grey or black.

27

28 Figure S4. Co-receptor preference of SHIV MNA.
29 SHIV MNA, along with controls for CCR5-tropic (SIV239) and CXCR4-tropic (HIV-1
30 NL4-3) and the parental HIV-1 MNA, were inoculated into TZM-bl cells in the
31 presence of increasing amounts of AD101 (Trkola *et al.*, 2002), provided by Dr. J.
32 Strizki, Schering Plough Research Institute, Kenilworth, NJ, and/or AMD3100
33 (Sigma-Aldrich, St. Louis, MO) (Donzella *et al.*, 1998).

34

35 Figure S5. Sensitivity of Env proteins from HIV-1 MNA and SHIV MNA to soluble
36 CD4 and small-molecule CD4 mimics.
37 Pseudotyped viruses carrying Env from SHIV MNA or HIV-1 MNA were normalised
38 by infectious titre at 100 TCID₅₀ and inoculated to TZM-bl cells in the presence of
39 increasing amounts of NBD-556, YYA-021 or soluble CD4. A pseudotyped virus
40 bearing A-MLV Env is acting as the negative control.

Figure S1




Figure S2




Figure S3




Figure S4




Figure S5

